

Effect of Oomycetes on Tree Declines: A Review

Muhannad Muhammad Nouri

Department of Biology, College of Education, Al-Iraqia University, Baghdad, IRAQ

Abstract

The influence of oomycetes can have significant impacts on Tree. Oomycetes have the most dangerous genus like *Phytophthora* and *Pythium*. These Oomycetes are genus of plant-damaging. *Phytophthora* can killed more than 1000 species of tree. This review discussed the role of oomycetes especially *Phytophthora* and *Pythium* on tree decline.

Keywords: urban; sub urban ; forest , Inciting factors, Predisposing factors

Received: 7 August 2022; **Revised:** 8 October 2022; **Accepted:** 15 October 2022; **Published:** 1 April 2023

1. Introduction

One of the most important contributing¹ to tree declines worldwide are diseases caused by pathogens in the genus *Oomycetes*. Once an invasive *Oomycetes* species arrives in a novel ecosystem it can cause significant losses to plant species and communities over large geographic areas. *Oomycetes ramorum* has destroyed huge areas of Californian forests²⁻⁴ and is responsible sudden oak death epidemic in the urban environment^{5,6}. *Oomycetes ramorum* is an introduced exotic pathogen to the USA, via the international nursery trade^{5,6}. Another introduced and invasive species is *P. cinnamomi*, it has an extensive host range of over 2000 plant species⁷, and contributes to White Oak Decline (*Quercus alba*) in the USA⁸. It has also destroyed huge areas of Australia's forests and woodlands⁹. For example, in the south-west of Western Australia, out of the 6510 described plant species, approximately 41% are susceptible to *P. cinnamomi*⁹. In Australia, *P. cinnamomi* is recognized as a 'Key threatening process to Australia's Biodiversity' by the Commonwealth Government's EPBC Act⁹, because of its direct impact on plant communities and indirect effects on native fauna through the loss of food, refugia, and habitat. For example, in Australia 157 *Banksia* species are recorded to be a major source of food for many kinds of an animal such as honeyeaters, rodents, honey possums, pygmy possums, gliders and bats^{10,11}. Many *Banksia* species are threatened by *Oomycetes* species^{9,12}. The impacts of *P. cinnamomi* in urban forests are poorly understood but likely to be significant in green space where it is present. The impact of *Oomycetes* diseases can have significant impacts on ecosystems, through declines in biodiversity¹³ and a decline in fundamental ecosystem services, such water equilibrium and river flow, modification of territorial climate patterns and carbon storage in biomass and soils. The impact of *Oomycetes* diseases can have significant impacts on ecosystems, through declines in biodiversity¹³ and a decline in fundamental ecosystem services, such water

equilibrium and river flow, modification of territorial climate patterns and carbon storage in biomass and soils^{14,15}.

2. The effect of predisposing and inciting factors on trees decline caused by Oomycetes species

Predisposing factors

Predisposing factors are climate or site factors that are long-term, static or slowly changing and render a tree vulnerable to a disease or disturbance^{16,17}. Trees damaged by these factors may slowly recover¹⁶ if further stresses are not imposed on the system. Predisposing factors play an important role in increasing the susceptibility of trees to *Oomycetes* species.

Water availability

Water is a key determining factor of forest health and includes waterlogging and flooding. Too little or too much water can predispose plants to pathogens, leading to forest diseases¹⁸. Waterlogging is the saturation of soil with water and can be exacerbated by poor drainage related to soil texture¹⁹ or in areas immediately adjacent to irrigation emitters²⁰. Waterlogging leads to low oxygen levels in soil, followed by anoxia or hypoxia and plant stress²¹. There are few tree species that can tolerate long-term low oxygen concentrations and even these species produce little canopy under heavy stress conditions²². Overall, soil saturation with water can predispose urban trees to infection by *Oomycetes* species. Corcobado *et al*²³ clearly showed how saturated soils were beneficial to *P. cinnamomi*, and disadvantageous to the tree leading to extensive root damage and death and decline of trees. Numerous other studies have shown tree declines linked with *Oomycetes* diseases are associated with waterlogging²⁴. In addition, excess water assists in the dispersal of *Oomycetes* species^{25,26}. Excess water through flooding can reduce the growth rate of trees in an urban area. How plants respond morphologically

and physiologically to extreme flooding is a significant part of determining the susceptibility of plants to pathogens²⁷.

Extreme weather

Recent climate models forecast greater fluctuations in a climate with increases in the frequency and severity of extreme events such as drought and heavy rains²⁸. The direct harm of extreme weather events on trees appears clearly through alterations in phenology, morphology, genetic frequencies and abundance of plants^{23,29}. Weather extremes will affect the growth and susceptibility of trees such as *Quercus ilex* to species such as *P. cinnamomi*²⁹. Also, extreme weather events could help pathogens to increase the number of their life cycles per year²⁹.

Extreme warming

Urban areas are generally warmer than adjacent rural areas, for example, the daily minimum temperature in New York City is on average ~7.2°F (~4°C) in summer months that makes it warmer than rural areas by an average 1.1°C³⁰. Elevated temperatures cause stress in trees and thus predisposes them to pathogens. Elevated temperature stress can have profound health consequences for plants through wilting, leaf burn, leaf folding and abscission, and physiological responses including changes in RNA metabolism and protein synthesis, enzymes, isoenzymes, and plant growth hormones. These changes will surely influence susceptibility to pathogens^{31,32}.

Inciting factors

Inciting factors are short-term physiological or biological factors that can increase the susceptibility of plants to pathogens by causing severe stress¹⁶. Trees influenced only by inciting factors may recover rapidly¹⁶. Indeed, the impact of inciting factors which include defoliation and excessive salt, on urban forest decline is a central challenge in forestry and ecology. Inciting factors could enhance the occurrence of trees decline by *Oomycetes* species.

Defoliation

Leaf defoliators have been deemed as causative factors of quick mortality in forest trees such as German oak trees³³. The link between *Oomycetes* species and defoliation was examined by^{34,35}. Jung *et al* Indicate that there is possibly a synergistic effect of *Oomycetes* spp. and defoliation events on plants³⁵ while Oszako *et al*³⁴ indicated defoliation can enhance the chances of *P. plurivora* to infect birch (*Betula pendula*).

Excessive salt

Salt can impact on tree growth by inhibiting chlorophyll synthesis through the direct effect on photochemistry reactions³⁶. Many studies reported that excessive salt enhances disease severity^{37,38}. These studies proved that soil salinity enhances *Oomycetes* Root Rot caused by *P. parasitica* in tomato³⁷ and stem rot caused by *P. citrophthora* in citrus rootstocks³⁸.

Loss of mycorrhizal fungi and other microorganisms

Mycorrhizal fungi are considered one of the most

important groups of microorganisms supporting the health of trees, as they help plants obtain nutrients and water through their mutualistic connection with roots of plants³⁹. Ectomycorrhizal fungi also reduce the severity of disease caused by *Oomycetes* species as they provide an effective barrier to *Oomycetes* infection. Watanarojanaporn *et al*⁴⁰ showed that root rot disease caused by *P. nicotianae* was reduced by arbuscular mycorrhiza. Similar results were found by Branzanti *et al*⁴¹ when investigating the correlation between the severity of *Q. robur* decline and mycorrhizal colonization and they showed that forest decline increased when mycorrhizal colonization decreased.

3. Discussion

Once introduced, urban landscapes can play an important role in the pathway of new *Oomycetes* species spreading into natural ecosystems. Many *Oomycetes* species have been recovered from urban environments. For example, 22% of type specimens of *Oomycetes* were isolated from urban environments and were described between 2001 and 2016⁴². The importance of nurseries for the dissemination of *Oomycetes* species cannot be underplayed as plant production nurseries supply material for urban and peri-urban plantings in addition to forestry, horticulture, and environmental restoration. Indeed, various *Oomycetes* species are considered as main pathogens of nursery plants⁴³. As the nursery trade continues to grow globally, *Oomycetes* species continue to spread into new areas, and nurseries are one of the principal routes of pathogen movement and introduction of alien plant pathogens into natural ecosystems⁴⁴. Hardy⁴⁴ found eight *Oomycetes* species associated with 14 nurseries in Perth, Western Australia. This indicates *Oomycetes* species have been problematic in nurseries for a long time in Perth.

4. Conclusion

The relationship between urban forest declines and *Oomycetes* species is considered important as they can result in economic and environmental losses including the loss of biodiversity and many tree species. There are many biotic and abiotic factors that can adversely impact on tree health leading to increased susceptibility of tree species to *Oomycetes*. However, there are gaps in our knowledge of the abiotic and biotic factors in urban or peri-urban landscapes that help predispose urban trees to *Oomycetes* diseases. This review has described the losses caused by *Oomycetes* species and has outlined some of the predisposing and inciting factors that can potentially influence the susceptibility of urban trees to *Oomycetes* species. It has also discussed some of the methods that can be used to rapidly identify *Oomycetes* species and to monitor their spread and impact across large areas, and how *Oomycetes* species can be managed.

Acknowledgment

The author would like to thank all people at the school of veterinary and life sciences Murdoch university, Australia.

References

- Orlikowski, L. B. *et al.* Oomycetes root and collar rot of mature *Fraxinus excelsior* in forest stands in Poland and Denmark. *For. Pathol.* 41, 510–519 (2011).
- Werres, S. *et al.* Oomycetes ramorum sp. nov., a new pathogen on *Rhododendron* and *Viburnum*. *Mycol. Res.* 105, 1155–1165 (2001).
- Rizzo, D. M. & Garbelotto, M. Sudden oak death: endangering California and Oregon forest ecosystems. *Front. Ecol. Environ.* 1, 197–204 (2003).
- Hansen, E. M., Parke, J. L. & Sutton, W. Susceptibility of Oregon forest trees and shrubs to Oomycetes ramorum: a comparison of artificial inoculation and natural infection. *Plant Dis.* 89, 63–70 (2005).
- Garbelotto, M., Rizzo, D. M. & Marais, L. Oomycetes ramorum and sudden oak death in California: IV. Preliminary studies on chemical control. *USDA For. Serv. Gen. Tech. Rep. No. PSW-GTR 184*, 811–818 (2002).
- Rizzo, D. M., Garbelotto, M., Davidson, J. M., Slaughter, G. W. & Koike, S. T. Oomycetes ramorum as the cause of extensive mortality of *Quercus* spp. and *Lithocarpus densiflorus* in California. *Plant Dis.* 86, 205–214 (2002).
- Hardham, A. R. Oomycetes cinnamomi. *Mol. Plant Pathol.* 6, 589–604 (2005).
- McConnell, M. E. & Balci, Y. Oomycetes cinnamomi as a contributor to white oak decline in mid-Atlantic United States forests. *Plant Dis.* 98, 319–327 (2014).
- Shearer, B. L., Crane, C. E. & Cochrane, A. Quantification of the susceptibility of the native flora of the South-West Botanical Province, Western Australia, to Oomycetes cinnamomi. *Aust. J. Bot.* 52, 435–443 (2004).
- Hackett, D. J. & Goldingay, R. L. Pollination of *Banksia* spp. by non-flying mammals in north-eastern New South Wales. *Aust. J. Bot.* 49, 637–644 (2001).
- Wrigley, J. W. & Fagg, M. Australian native plants: cultivation, use in landscaping and propagation. (*No Title*) (2013).
- McCredie, T. A., Dixon, K. W. & Sivasithamparam, K. Variability in the resistance of *Banksia* Lf species to Oomycetes cinnamomi Rands. *Aust. J. Bot.* 33, 629–637 (1985).
- Garkaklis, M. J., Calver, M. C., Wilson, B. A. & Hardy, G. E. S. J. Habitat alteration caused by an introduced plant disease, Oomycetes cinnamomi: a significant threat to the conservation of Australian forest fauna. in *Conservation of Australia's Forest Fauna*, 2nd edition 899–913 (Royal Zoological Society of New South Wales, 2004).
- Foley, J. A. *et al.* Amazonia revealed: forest degradation and loss of ecosystem goods and services in the Amazon Basin. *Front. Ecol. Environ.* 5, 25–32 (2007).
- Alford, R. A. & Richards, S. J. Global amphibian declines: a problem in applied ecology. *Annu. Rev. Ecol. Syst.* 30, 133–165 (1999).
- Worrall, J. J. *et al.* Effects and etiology of sudden aspen decline in southwestern Colorado, USA. *For. Ecol. Manage.* 260, 638–648 (2010).
- Aghighi, S. *et al.* A conceptual model to describe the decline of European blackberry (*Rubus anglocandicans*), a weed of national significance in Australia. *Plant Dis.* 98, 580–589 (2014).
- Doblas-Miranda, E. *et al.* A review of the combination among global change factors in forests, shrublands and pastures of the Mediterranean Region: Beyond drought effects. *Glob. Planet. Change* 148, 42–54 (2017).
- Bryla, D. R. & Linderman, R. G. Implications of irrigation method and amount of water application on Oomycetes and Pythium infection and severity of root rot in highbush blueberry. *HortScience* 42, 1463–1467 (2007).
- Café-Filho, A. C. & Duniway, J. M. Effect of location of drip irrigation emitters and position of Oomycetes capsici infections in roots on Oomycetes root rot of pepper. *Phytopathology* 86, 1364–1369 (1996).
- Lambers, H., Chapin, F. S. & Pons, T. L. *Plant physiological ecology*. vol. 2 (Springer, 2008).
- Kozłowski, T. T., Kramer, P. J. & Pallardy, S. G. Physiological ecology the Physiological Ecology of Woody Plants. at (1991).
- Corcobado, T., Cubera, E., Moreno, G. & Solla, A. *Quercus ilex* forests are influenced by annual variations in water table, soil water deficit and fine root loss caused by Oomycetes cinnamomi. *Agric. For. Meteorol.* 169, 92–99 (2013).
- Shearer, B. L. & Fairman, R. G. A stem injection of phosphite protects *Banksia* species and *Eucalyptus marginata* from Oomycetes cinnamomi for at least four years. *Australas. Plant Pathol.* 36, 78–86 (2007).
- Baker, K. F. & Matkin, O. A. Detection and control of pathogens in water. *Ornamentals Northwest Newsl.* 2, 2–14 (1978).
- Stanghellini, M. E., Rasmussen, S. L., Kim, D. H. & Rorabaugh, P. A. Efficacy of nonionic surfactants in the control of zoospore spread of *Pythium aphanidermatum* in a recirculating hydroponic system. *Plant Dis.* 80, 422–428 (1996).
- Stolzy, L. H. & Sojka, R. E. Effects of flooding on plant disease. *Flooding and plant growth* 221–264 (1984).
- Solomon, S. *Climate change 2007-the physical science basis: Working group I contribution to the*

- fourth assessment report of the IPCC*. vol. 4 (Cambridge university press, 2007).
29. Corcobado, T., Cubera, E., Juárez, E., Moreno, G. & Solla, A. Drought events determine performance of *Quercus ilex* seedlings and increase their susceptibility to *Oomycetes cinnamomi*. *Agric. For. Meteorol.* 192, 1–8 (2014).
 30. Rosenzweig, C., Solecki, W. & Slosberg, R. Mitigating New York City's heat island with urban forestry, living roofs, and light surfaces. *A Rep. to New York State Energy Res. Dev. Auth.* 1–5 (2006).
 31. Chuine, I., Cambon, G. & Comtois, P. Scaling phenology from the local to the regional level: advances from species-specific phenological models. *Glob. Chang. Biol.* 6, 943–952 (2000).
 32. Garrett, K. A., Dendy, S. P., Frank, E. E., Rouse, M. N. & Travers, S. E. Climate change effects on plant disease: genomes to ecosystems. *Annu. Rev. Phytopathol.* 44, 489–509 (2006).
 33. Thomas, F. M., Blank, R. & Hartmann, G. Abiotic and biotic factors and their interactions as causes of oak decline in Central Europe. *For. Pathol.* 32, 277–307 (2002).
 34. Oszako, T. *et al.* Assessment of interactions between defoliation and *Oomycetes plurivora* stem infections of birch seedlings. *For. Chron.* 94, 140–146 (2018).
 35. Jung, T., Blaschke, H. & Oßwald, W. Involvement of soilborne *Oomycetes* species in Central European oak decline and the effect of site factors on the disease. *Plant Pathol.* 49, 706–718 (2000).
 36. Parida, A. K., Das, A. B. & Mittra, B. Effects of salt on growth, ion accumulation, photosynthesis and leaf anatomy of the mangrove, *Bruguiera parviflora*. *Trees* 18, 167–174 (2004).
 37. Swiecki, T. J. & MacDonald, J. D. Soil salinity enhances *oomycetes* root rot of tomato but hinders asexual reproduction by *Oomycetes* parasitic. *J. Am. Soc. Hortic. Sci.* 116, 471–477 (1991).
 38. Sulistyowati, L. & Keane, P. J. Effect of soil salinity and water content on stem rot caused by *Oomycetes citrophthora* and accumulation of phytoalexin in citrus rootstocks. *Phytopathology* 82, 771–777 (1992).
 39. Sapsford, S. J., Paap, T., Hardy, G. E. S. J. & Burgess, T. I. The 'chicken or the egg': which comes first, forest tree decline or loss of mycorrhizae? *Plant Ecol.* 218, 1093–1106 (2017).
 40. Watanarojanaporn, N., Boonkerd, N., Wongkaew, S., Prommanop, P. & Teaumroong, N. Selection of arbuscular mycorrhizal fungi for citrus growth promotion and *Oomycetes* suppression. *Sci. Hortic. (Amsterdam)*. 128, 423–433 (2011).
 41. Branzanti, M. B., Rocca, E. & Pisi, A. Effect of ectomycorrhizal fungi on chestnut ink disease. *Mycorrhiza* 9, 103–109 (1999).
 42. Hulbert, J. M., Agne, M. C., Burgess, T. I., Roets, F. & Wingfield, M. J. Urban environments provide opportunities for early detections of *Oomycetes* invasions. *Biol. Invasions* 19, 3629–3644 (2017).
 43. Erwin, D. C. & Ribeiro, O. K. *Oomycetes diseases worldwide*. (American Phytopathological Society (APS Press), 1996).
 44. Hardy, G. E. & Sivasithamparam, K. *Oomycetes* spp. associated with container-grown plants in nurseries in Western Australia. *Plant Dis.* 72, 435–437 (1988).