

# Structural Studies of Valine Complexes with Biological Activity

Athir A.R. Mahmud

College of Medicine, Al-Iraqia University, Al-Aadhamiya District, Baghdad, IRAQ

## Abstract

A transition metal ion combination of Ni(II) ion with mixed ligands has been produced, whereby Valine serves as the secondary ligand and 1,10-phenanthroline (phen) in its principal role. The ability of the complex to stop the growth diameter of three different species of fungi—*Aspergillus flavus*, *Aspergillus Niger*, and *Pencillium*—was used to measure its antifungal properties at different concentrations.

**Keywords:** Valine complexes; Antifungal activity; Structural Studies; Biological activity

**Received:** 2 August 2024; **Revised:** 3 September 2024; **Accepted:** 10 September 2024; **Published:** 1 October 2024

## 1. Introduction

With the advancement in the field of bioinorganic chemistry the role of transition metal complexes as therapeutic compounds is becoming increasingly important. The use of transition metal complexes as therapeutic compounds has become more and more pronounced [1]. Metal ions play an important role in bioinorganic chemistry and metals such as Fe, Co, Ni, Cu, Zn, and Cd may exist in trace amounts in biological systems. Structural studies of the complexes of these metals with biological compounds are extremely important [2]. The study of the coordinated systems metal ion - amino acids has become increasingly important in recently from different points of view. Increasing worldwide interest was confirmed in 1991, when EC countries selected "Biocoordination Chemistry" as one of seven priorities research fields [3]. The ligand 1,10-phenanthroline (phen) as in figure (1) is a strong field bidentate ligand that forms very stable chelates with first row elements of transition metals [4].




Figure (1) structure of 1,10-phenanthroline

Although a good number of metal mixed-ligand complexes containing either 2,2'-bipyridine or 1,10-phenanthroline with some other ligands are reported in the literature, there are a few reports on the synthesis and biological activities of metal mixed-ligand complexes containing 2,2'-bipyridine and 1,10-phenanthroline as ligands [5].

1,10-phenanthroline (phen) whether in the metal-free state and coordinated to the transition metals disturb the functioning of a wide variety of biological system[6].

Valine is an  $\alpha$ -amino acid with the chemical formula  $\text{HO}_2\text{CCH}(\text{NH}_2)\text{CH}(\text{CH}_3)_2$  as in Fig. (2). L-Valine is one of 20 proteinogenic amino acids. This essential amino acid is classified as non-polar. Valine is branched-chain amino acid. Human dietary sources include cottage cheese, fish, poultry, peanuts, sesame seeds, and lentils [7]. Valine has many properties [8-9] shows in table (2).




Figure (2) structure of Valine

Amino acids are organic molecules that constitute a most important part of biological structure and

body chemistry. In the amino acids, the amino group is basic portion of the molecule and it is capable of reacting with both organic and inorganic acids to form salts, while the acid portion is capable of reacting with bases [10]. L-Valine complexes belong to non-centrosymmetric space groups and it is an essential criterion for nonlinear optical (NLO) application. L-Valine hydrogen chloride (LVHC) crystal has been formed by combining L-Valine and hydrochloric acid and this complex is found to be a useful NLO material. It has been reported that doping NLO crystals with organic and inorganic impurities can alter various physical and chemical properties and doped-NLO crystals may also find applications in optoelectronic devices like pure NLO crystals [11-14].

*Aspergillus Niger* or *A. Niger* is a fungus and one of the most common species of the genus *Aspergillus*. It causes a disease called black mold on certain fruits and vegetables such as grapes, onions, and peanuts, and is a common contaminant of food. It is ubiquitous in soil and is commonly reported from indoor environments, where its black colonies can be confused with those of *Stachybotrys* (species of which have also been called "black mould") [15]. Some strains of *A. Niger* have been reported to produce potent mycotoxins called ochratoxins [16], but other sources disagree, claiming this report is based upon misidentification of the fungal species. Recent evidence suggests some true *A. Niger* strains do produce ochratoxin A. [17].

*Aspergillus flavus* is a saprotrophic and pathogenic [18] fungus with a cosmopolitan distribution [19]. *A. flavus* infections can occur while hosts are still in the field (pre-harvest), but often show no symptoms (dormancy) until post-harvest storage and/or transport. In addition to causing pre-harvest and post-harvest infections, many strains produce significant quantities of toxic compounds known as mycotoxins, which when consumed are toxic to mammals [20].

Species of *Penicillium* are ubiquitous soil fungi preferring cool and moderate climates, commonly present wherever organic material is available. Saprophytic species of *Penicillium* and *Aspergillus* are among the best-known representatives of the Eurotiales and live mainly on organic biodegradable substances. Commonly known as molds, they are among the main causes of food spoilage, especially species of subgenus *Penicillium* [21]. Many species produce highly toxic mycotoxins. The ability of these *Penicillium* species to grow on seeds and other stored foods depends on their propensity to thrive in low humidity and to colonize rapidly by aerial dispersion while the seeds are sufficiently moist [22].

## 2. Materials and Methods

### Preparation of complex

Complex was prepared in general method of mixed ligands metal complexes [23-24] with preparation a solution of 1,10-phenanthroline (0.180 g, 1 m.mol) in aqueous ethanol (1:1:10 ml) and solution of L-Valine (0.234, 2 m.mol) in aqueous ethanol (1:1:10 ml) containing sodium hydroxide (0.08, 2mmol) were added simultaneously to a solution of  $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$  (1 m.mol) in aqueous ethanol (1:1:10 ml) in the stoichiometric ratio [2Val:M: phen] as (Scheme 1). The above solution was stirred for 1 hour and allowed to stand for overnight. The product formed was filtered off, washed with aqueous ethanol (1:1 ml) and dried in air.



### Antifungal activity test

Fungi were obtained from the University of Mustansiriyah and diagnosed based on the taxonomic keys in Domch & Gamas. The antifungal activity was determined on the potato dextrose agar which it was solidified. According to the manufacturer's instructions Attended by dissolving 39 g of powdered middle-ready and as directed by the manufacturer in 1000 ml of distilled water and acidic function seized to 5.6 and infertility medium, added 1ml of prepared complex and added the media after let go solidifies. The percentage of inhibition was measured by the following equation:

$$\text{percentage of inhibition} = \frac{\text{The growth of fungus on the control medium} - \text{The growth of fungus on the Medium with complex}}{\text{The growth of fungus on the control media}} \times 100$$

## 3. Results and Discussion

Generally, the complexes were prepared by reacting the respective metal salts with the ligands using 1:1:2 mole ratio, i.e. one mole of metal salt: one mole of 1,10-phenanthroline and two moles of sodium valinate. The synthesis of mixed ligand Metal complexes may be represented as follows



Where phen is 1,10-phenanthroline and Val H is amino acid L-valine).

The formula weight, molecular weight, color and melting points are given in (Table 3). Based on the physicochemical characteristics (Table 3), it was found that all the complexes were non-hygroscopic, stable at room temperature.

The results of the antifungal activity of different concentrations of complex  $[\text{Ni}(\text{Val})_2(\text{Phen})]$  on the radial growth of plant pathogenic fungi compared with the control, the concentration 0.01M is the best among the concentrations, where it was diameter in cm unit of growth *Aspergillus Niger* a solid medium 1.5cm and

*Pencillium* 2.6cm while a growth *Aspergillus flavus* 1.2cm while the concentration 0.001M was ranged affect between weak and had to growth of *Aspergillus Niger* as reached diameter of growth 2.5cm, and *Pencillium* 3.8cm to a high affect on *Aspergillus flavus* as growth zone was 2.6cm compared with the control media in growth of three fungi, finally the concentration 0.0001M was absent in affect on *Pencillium* in growth zone was 4cm Equal with control medium and semi-absent affect on *Aspergillus Niger* in growth zone was 3.8cm Comparison with control medium which stood at 4cm while affect on *Aspergillus flavus* was highest and it was growth zone 3.4cm Comparison with control medium which stood at 4cm .

Although the *Aspergillus flavus* was the best among the three fungi in growth in the control media, but it was most sensitive direction of the chemical complex. The results were viewed in tables (4,5,6). The results of affect the different concentrations of complex on plant pathogenic fungi show below in Fig. (1).




Fig. (1) affects the different concentrations of complex on types of fungi

## References

- [1] Eshwika et al, (2004), Metal complexes of 1, 10-phenanthroline-5,6-dione alter the susceptibility of the yeast *Candida albicans* to Amphotericin B and Miconazole, in *Bio metals*, Vol.17,p. 415-422 .
- [2] Canpolat et al, (2004), synthesis, characterization of some Co(II) complexes with Vic-Dioxime ligands and their antimicrobial properties. *Turk. j. chem.*28, 235-242.
- [3] Yuichi Shimazaki, Masako Takanib and Osamu Yamauchi, Metal complexes of amino acids and amino acid side chain groups. *Structures and properties Dalton Trans.*,7854–7869(2009).
- [4] Lee, J.D concise *Inorganic chemistry*, 4th ed., Chapman and Hill: London; (1991); pp 607-1074.
- [5] Hazel, A.; Simonsen, O.; Wernberg, O. *Acta. Cryst.* 1986, C42, 1707.
- [6] Deegan et al. (2007).
- [7] Nomenclature and symbolism for amino acids and peptides (IUPAC-IUB Recommendations, *Pure Appl. Chem.* 56(5): 595 (1984).
- [8] Dawson, R.M.C., et al., *Data for Biochemical Research*, Oxford, Clarendon Press, (1959).
- [9] Weast, Robert C., ed. (1981). *CRC Handbook of Chemistry and Physics* (62nd ed.). Boca Raton, FL: CRC Press. p. C-569. ISBN 0-8493-0462-8.
- [10] Venkata et al. *Bull.chem. soc. Ethiop.* (2009). Vol.23,No.2, p.197-204 .
- [11] C. Krishnan, P. Selvarajan, T.H. Freeda (2008). *J. Crystal Growth* 311:141.
- [12] S. Goma, C.M. Padma, C.K. Mahadevan (2006) *Materials Letters* (60):3701.
- [13] P. Selvarajan, A. Siva dhas, T.H. Freeda, C.K. Mahadevan (2008). *Physica B* 403: 205.
- [14] N.P. Rajesh, V. Kannan, P.S. Raghavan, P. Ramasamy, C.W. Lan (2002). *Mater. Lett.* (52):326.
- [15] Samson RA, Houbraken J, Summerbell RC, Flannigan B, Miller JD (2001). Common and important species of fungi and actinomycetes in indoor environments. In: *Microorganisms in Home and Indoor Work Environments*. New York: Taylor & Francis. pp. 287–292. ISBN.
- [16] Abarca M, Bragulat M, Castellá G, Cabañes F (1994). "Ochratoxin A production by strains of *Aspergillus Niger* var. *Niger*". *Appl. Environ Microbiol* 60 (7): 2650 .
- [17] Schuster E, Dunn-Coleman N, Frisvad JC, Van Dijck PW (August 2002). "On the safety of *Aspergillus Niger* - a review". *Applied microbiology and biotechnology* 59 (4–5): 426–35.
- [18] Masayuki Machida; Katsuyai Gomi (2010). *Aspergillus: molecular biology and genomics*. Horizon Scientific Press. pp. 157.
- [19] Ramírez-Camejo, L. A.; Zuluaga-Montero, A.; Lázaro-Escudero, M. A.; Hernández-Kendall, V. N.; Bayman, P. (2012). "Phylogeography of the cosmopolitan fungus *Aspergillus flavus*: Is everything everywhere?". *Fungal Biology* 116 (3): 452–463.
- [20] Agrios, George N. (2005). *Plant Pathology: Fifth Edition*. Elsevier Academic Press. p. 922.

- [21] Samson RA, Seifert KA, Kuijpers AF, Houbraken JA, Frisvad JC. (2004). "Phylogenetic analysis of *Penicillium* subgenus *Penicillium* using partial beta-tubulin sequences" (PDF). *Studies in Mycology* 49: 175–200.
- [22] Pitt JI, Basílico JC, Abarca ML, López C. (2000). "Mycotoxins and toxigenic fungi". *Medical Mycology* 38(Suppl 1): 41–46.
- [23] P.K. Bhattacharya, H.J. Lawson, J.K. Barton, (1H NMR Studies of Nickel(II) Complexes Bound to Oligonucleotides: A Novel Technique for Distinguishing the Binding Locations of Metal Complexes in DNA) *Inorg. Chem.* 42, 8811-8817 (2003).
- [24] K. Brugger, "Coordination Chemistry Experimental Methods", London butterworth's England (1973).

**Table (1) Physical properties of 1,10-phenanthroline**

Compound	Molecular Formula	Physical State	Molar mass (g/mol)	Solubility (in water)	Acidity (pKa)	Melting Point (°C)
1,10-phenanthroline	C <sub>12</sub> H <sub>8</sub> N <sub>2</sub>	White Crystalline Powder	180.21	Slightly Soluble	4.27	114-120

**Table (2) Physical properties of Valine**

Compound	Molecular Formula	Density (g/cm <sup>3</sup> )	Molar mass (g/mol)	Solubility (in water)	Acidity (pKa)	Melting Point (°C)
Valine	C <sub>5</sub> H <sub>11</sub> NO <sub>2</sub>	1.316	117.15	Soluble	2.32 (carboxyl) 9.62 (amino)	298 Decomp.

**Table (3) Physical properties of complex**

Compound	Molecular Weight	Color	Melting Point (°C)	% Metal Theory
[Ni(phen)(Val) <sub>2</sub> ]	471.18	Light green	273	12.46

**Table (4) Antifungal activity of different concentration of complex on radial growth rate of *Aspergillus Niger*, *Aspergillus flavus*, *Penicillium sp.***

Fungi	Control cm	Conc. of complex Mol/lit	Diameter of growth Cm	% Inhibition
<i>Aspergillus Niger</i>	4	0.01	1.5	62.5
		0.001	3.5	12.5
		0.0001	3.8	5
<i>Aspergillus flavus</i>	5	0.01	1.2	76
		0.001	2.6	48
		0.0001	3.4	32
<i>Penicillium sp.</i>	6	0.01	2.6	35
		0.001	3.8	5
		0.0001	4	0